



INSTITUTE FOR
MARINE RESEARCH
DAUIN · PHILIPPINES

POBLACION REEF STATUS REPORT

2019





INSTITUTE FOR
MARINE RESEARCH
DAUIN · PHILIPPINES

POBLACION REEF

STATUS REPORT

2019

DISCLAIMER

The research reported herein is based on initial analyses of complex datasets as part of the Dauin Reef Long Term Monitoring Project, and should not be considered definitive in all cases. Institutions or individuals interested in the results or applications of the Institute for Marine Research are invited to contact the Director at the Dauin address below.

For additional copies of this report, please phone IMR on (+63) 917 110 0421 or write to us at info@institutemarinereserch.org

This report, along with a range of information about IMR, is available online at www.institutemarineresearch.org

© Institute for Marine Research

Institute for Marine Research
Tugawe, Municipality of Dauin
6217, The Philippines

Front Cover: Poblacion Marine Reserve, Dauin, The Philippines. Image: Tracey Jennings

Full report written by:

Chelsea Waters
Director, Institute for Marine Research
chelsea@institutemarineresearch.org

EXECUTIVE SUMMARY

The world's coral reefs are being severely degraded by the activities of humans, and the need to reduce local threats to offset the effects of increasing global pressures is now widely recognized. Major anthropogenic risk factors include mortality and reduced growth of reef-building corals due to their high sensitivity to rising seawater temperatures, ocean acidification, deteriorating water quality, destructive fishing, over-exploitation of key marine species, and the direct devastation of coastal ecosystems through unsustainable coastal development^{23,41}. These anthropogenic risks interact with other large-scale acute disturbances, including tropical storms and population outbreaks of the corallivorous crown-of-thorns starfish (COTS) *Acanthaster planci*, which may also increase in frequency and intensity in response to human activities. Regional policies can no longer protect reefs from global-scale devastation due to climate change-associated heat stress and intensifying tropical storms⁴¹. Efforts are therefore shifting toward management of local and regional anthropogenic pressures to strengthen reef resilience. However, assessment of the likely effectiveness of reductions of local anthropogenic pressures requires a sound understanding of the processes that determine ecosystem trajectories.

The Philippines, represents a particularly relevant case to investigate ecosystem trajectories. Over 7,100 islands dominate the Philippine archipelago, which is located within the heart of the incredible biological diversity that is the 'Coral Triangle'. Boasting 76% of the world's total coral species and 37% of the reef fishes of the world⁴², this incredible biological diversity of the Coral Triangle is associated with some of the highest human population densities and growth rates in the world²³. Changes to the health of coastal ecosystems are exposing

coastal populations to the erosion of food security and income, deteriorating coastal protection and other challenges. They are affecting people who are already impoverished and are among the least able to respond to the changes that are occurring in their environment²³. Reef fisheries have estimated to directly contribute to 15 – 30% of the Philippines total known national municipal fisheries (obtained from licenses issued through local-government areas), with nearly 70% of the protein food intake being fish. The stark contrast between poverty, hunger and deprivation amidst this increasing want is rapidly declining reef resources. It is therefore no surprise that it is in the Philippines that reefs are at the highest risk from overexploitation, destructive fishing and other human related impacts such as coastal development and sedimentation. If these processes are allowed to continue, these changes will exacerbate poverty and social instability within the region, with wider consequences for the region and the world. It is imperative that we address the core issue of anthropogenic climate change whilst at the same time addressing the key threats that are rising from local stressors.

ABBREVIATIONS

ABBREVIATION	TERM IN FULL
1-D	Simpsons Index of Diversity
2D	2-Dimensional
3D	3-Dimensional
AIMS	Australian Institute of Marine Science
BBD	Black Band Disease
BrBD	Brown Band Disease
CPC _e	Coral Point Count with Excel Extension
COTS	Crown of Thorns Starfish
DEM	Digital Elevation Model
HYP	Hyperplasia
IMR	Institute for Marine Research
LTRMP	Long Term Reef Monitoring Project
NEO	Neoplasia
PP	Porites Pinking
S	Species Richness
SCUBA	Self Contained Underwater Breathing Apparatus
SE	Standard Error
SEB	Skeletal Eroding Band
SfM	Structure from Motion
SVS	Stereo Video System
UVC	Underwater Visual Census

TABLE OF CONTENTS

1. INSTITUTE FOR MARINE RESEARCH	7
2. A MESSAGE FROM THE DIRECTORS	8
3. METHODOLOGY	9
3.1 SURVEY SITES	9
3.2 RESEARCH TECHNIQUES	10
3.2.1 3-DIMENSIONAL REEF MODELLING	10
3.2.2 DIVER OPERATED STEREO VIDEO SYSTEM	11
3.2.3 BENTHIC ASSAYS	11
3.2.4 SCUBA SEARCH	12
4. RESULTS	13
4.1 BENTHIC COVER	13
4.2 REEF IMPACTS & CORAL MORTALITY	13
4.3 FISH	14
4.4 REEF COMPLEXITY	14
5. DISCUSSION	16
6. REFERENCES	18

1. INSTITUTE FOR MARINE RESEARCH

The Institute for Marine Research (IMR) is a grassroots non-profit organization that conducts long-term and fine-scale research on coastal marine ecosystems, using this scientific evidence to educate, transform and encourage locally led marine conservation strategies within the Municipality of Dauin.

The Institute will deliver the science to help realize three key long-term impacts for the Municipality:

1. Improve the health and resilience of marine and coastal ecosystems across the Municipality
2. Ensure economic, social and environmental net benefits for Dauin's marine industries and coastal community
3. Protect Dauin's coral reefs and other tropical marine environments from the effects of climate change and coastal development

The Dauin *Long-Term Reef Monitoring Project (LTRMP)* was established by IMR in February 2019 to track fine-scale changes in the overall reef community of Dauin's fringing reef system, and realize the three key long-term impacts for the Municipality. More specifically, the aims of the Dauin *LTRMP*:

1. Understand how benthic composition (measured as percentage cover, species diversity indices, species

abundance, structural complexity, slope and rugosity) influences fish community structure (measured through biomass, species abundance, trophic groups, and species diversity indices)

2. Document the effect of disturbances such as *Acanthaster planci* (Crown of Thorns Starfish, COTS) and *Drupella* spp. outbreaks, typhoons, and bleaching events. The data will also provide awareness of other threats to the reef (such as coral disease, human activity, illegal poaching, high nutrient outflow, trash) that will be of concern to reef managers
3. Document the effects of temperature, changing light regimes, dissolved oxygen, and pH on the seasonal and annual variability of Dauin's fringing reef

All results collected as part of the *LTRMP* will be used to:

- a) Publish and present annual Outlook reports to policy-makers within the Local Government Unit (LGU)
- b) Determine 'areas of concern' with regards to unsustainable practices occurring within the Municipality
- c) Publish findings on a wider scientific platform to expand our current knowledge of coral reef ecosystems

2. A MESSAGE FROM THE DIRECTORS



What an action-packed and rewarding start to our first research season here in the Philippines! With 19 research sites within the Municipality of Dauin, we have this reef system well monitored!

With that being said, these results are just the beginning.

We have a long road to go with deepening our research to understand the resiliency state of our reef system towards the threats and challenges associated with our changing climate. On a localised platform, our results are catching a glimpse of the negative, human-induced practices that are exacerbating coral mortality within the region.



Our first step towards reef conservation is awareness and partnership. We are proudly partnering not only with Dauins Local Government Unit (LGU), but with various local resorts, NGOs and other local stakeholders who wish to share our common goal of preserving Dauins coastal ecosystem.

We would also like to take this opportunity to say how proud and thankful we are of our Research Assistants and Fellows who have not only assisted the Institute in meeting our seasonal research objectives, but for everything that comes both afterwards and in-between. From the months of data analysis, to the weeks of interpretation of findings into site reports. From creating school lesson plans, and environmental awareness initiatives. You have helped to take IMR to a whole new level. Our heartfelt thanks to you all.

- Chelsea Waters & Rafael Manrique

3. METHODOLOGY

3.1 SURVEY SITES

Dauin is a fourth class Municipality in the province of Negros Oriental, Philippines. The Municipality stretches across nine kilometres of coastline, bordered in the north by Bacong, and Zamboanguita in the south. Nineteen core sites were selected for seasonal and annual monitoring. These sites span the variation in the coral reef composition of benthic and fish communities across the Municipality, and account for the zoning history of its associated no-take marine protected areas. The nineteen core sites consist of fifty metre transects that are laid out parallel to the reef crest, between

depth ranges of 1 – 6 metres and 7 – 12 metres. Surveys are conducted bi-annually to account for seasonal variability, with “dry” season surveys running from February to July, and “wet” season surveys running from August to January.

Poblacion, located within Barangay Poblacion I. Four fifty metre replicates (n = 4) were conducted between the months of February and July.




Figure 3.1. Location of the Municipality of Dauin and IMRs survey sites on Negros Oriental, the Philippines. Maps sourced from GADM database of Global Administrative Areas (2015) under a CC BY licence, used with permission.

3.2 RESEARCH TECHNIQUES

3.2.1 3-Dimensional Reef Modelling

A 3D camera rig consisting of two *GoPro Hero 5 Black* cameras attached to a one-metre long aluminium pole is assembled. The cameras are placed 90 centimetres apart, having one on each end of the pole⁴⁷. The cameras are placed in a downward facing position at the beginning of the 50 metres. The aim of the diver is to get over 60% overlap from

pictures to ensure they can be aligned, with preliminary testing indicating this method decreases alignment errors over single passes or higher image intervals³². The rig is kept approximately 2 meters above the substrate with the cameras always aimed straight down at the substratum³⁵. A lawnmower pattern is conducted at a steady pace, 1 metre either side of the transect.

Introduction to 3-Dimensional Reef Modelling

Structural complexity is a key habitat feature that influences ecological processes by providing a set of primary and secondary resources to organisms, such as shelter from predators and availability of food. The spatial configuration and morphology of corals create complex structures that serve as habitats for a large number of species inhabiting coral reefs. As such, structural complexity of coral reefs drives numerous functions directly linked to the resilience of these ecosystems^{45,46}.

Despite the importance of reef structure in the long-term functioning of these systems, quantifying its complexity is a time-consuming exercise. Therefore, advancing our understanding of how structural complexity influences reef dynamics requires improving our efficiency and ability to quantify multiple metrics of 3D structural complexity in a repeatable way, across spatial extents, whilst maintaining a high resolution.

IMR researchers are making use of rapid advances in technology to monitor reef structural complexity by recreating and measuring reefs in 3D. Using off-the-shelf cameras, the 3D structure of the reef is accurately reconstructed by underwater images taken at pace across a reef transect. These images are aligned and referenced using a technique called photogrammetry, which allows the recovery of the exact position of each pixel in the images, recreating the 3D structure of the reef^{43,44}.

These 3D models are produced at scale, allowing IMR scientists to measure different attributes associated with the structural complexity of coral reefs, such as surface complexity (3D/2D surface area), curvature, volume and slope, across large extents in a fraction of the time that takes to do it underwater. With the advances in photogrammetry software and high performance computing hardware, automated analyses of structural complexity across all IMR-monitored reefs in Dauin is now possible and at a minimal cost. Characteristics of the reef surface are believed to play an important part in the early life of corals and subsequent reef recovery. We can now measure things we could never measure before, including being able to see how complex the surface of the reef is.

3.2.2 Diver Operated Stereo Video System

Transects are conducted using a diver-operated Stereo-Video System (SVS; SeaGIS, Melbourne, Australia), comprised of two GoPro Hero 5 Black video cameras. Transects are 50 metres long following the reef contour. Surveys are conducted by two people; the SVS operator and a second diver responsible for distance measurements. To minimise potential disturbance to the fish community, cameras are set to record and synchronised prior to entry. Transects begin with the cameras pointing vertically down, until alerted via a fin tug to indicate the start of the transect. At this point cameras are now pointed along the reef, with another fin tug indicated the end of the transect after a further 50 metres.

Cameras are angled approximately 20° downwards, and kept approximately 0.5 metres above the substrate, filming the reef scape along the transect. Transects take approximately 5 - 6 minutes to film using SCUBA. Footage is analysed in EventMeasure software v3.51 (SeaGIS, Melbourne, Australia) allowing the calibrated SVS footage to be synchronised and fish total lengths to be measured. EventMeasure also resolves centre points of each individual fish encountered into distances on a three-dimensional coordinate system, allowing the exclusion of fish outside 2.5 metre either side and 5 metre in front of the camera system. Side distance restrictions maintains a consistent belt along the transect, while a front distance restriction prevents variations in visibility (e.g. turbidity, light intensity) from influencing data.

Introduction to the Diver Operated Stereo Video System

Understanding of fish ecology, and our ability to effectively manage fish populations requires accurate data on diversity, abundance and size. Underwater visual census (UVC) surveys have been a widely used method to collect data on coastal fish assemblages. UVC requires divers to identify and count fishes within a predetermined area, or by distance-based sampling. This is a logistically simple, non-destructive, and cost-effective method of surveying fish. However, the effectiveness of UVC for reliable long-term monitoring is influenced by inter-observer variability and inaccuracies in estimating the length of fish and sampling areas. In addition, a combination of the identification, counting and size estimations of fish requires extensive training and experience.

IMR utilises a Diver Operated Stereo Video System, an innovative technology which allows our researchers to not only record fish species with more precision and accuracy than the traditional Underwater Visual Census (UVC) techniques, but efficiently quantifying the abundance and size of reef fish^{12,48}. Rather than relying on in situ identification and length estimates, collected video data can be annotated in the lab reducing time in the field and/or enabling greater spatial coverage.

3.2.3 Benthic Assays

Benthic surveys of stationary benthic organisms are conducted following the technique of the Australian Institute of

Marine Science (AIMS) LTMP. Benthic surveys are conducted along the transect line. At each site, single frames are shot at 1 metre intervals using a GoPro camera. Fifty still frames are shot along each 50 metre transect, with the camera held

approximately 50 centimetres above the substrate. Photographs are analysed through the use of CPCe software by Kohler and Gill (2006). Underwater photographic frames are overlaid by a matrix of randomly distributed points. In this case, thirty random points are overlaid and generated in the whole frame of each photo and used for identification. Point overlay is used to characterise the benthos, and determine percentage type of organism and substrate in the image⁴⁹. The species code data for each frame is

stored in a .cpc file which contains the image filename, point coordinates and the identified data codes. The data from individual frames can be combined to produce inter and intra transect and site comparisons via automatically generated Excel spreadsheets. For each category of benthic organism, the mean values for percent cover at each site are used to estimate seasonal and temporal trends in cover of benthic organisms at each site, zone, and throughout the municipality as a whole.

Introduction to Benthic Assays

With the world's coral reefs being severely degraded by the activities of humans, there is a need to efficiently assess and monitor reefs even at the regional and local level^{51,52}. Coral Point Count (CPCe) is a visual basic software designed to quickly and efficiently calculate statistical coral coverage over a specified area through the aid of photo-transects⁵⁰. These transect images are assigned with spatial random points for user's further identification. It can also perform both image calibration and area analysis of the benthic features, and has the ability to automatically generate analysis in Microsoft Excel. Thus, CPCe is a highly significant useful tool, particularly in coral reef monitoring, assessment and conservation.

3.2.4 SCUBA Search

SCUBA searches are designed to provide a more detailed picture of the causes and relative scale of coral mortality, and are conducted following a modified version of AIMS LTMP. SCUBA searches are made along a fixed 50 m transect, with a 2 m belt (1 metre either side of the central tape measure). Numbers are recorded for

the following: crown-of-thorns starfish (COTS), COTS feeding scars, *Drupella* spp., *Drupella* spp. feeding scars, unknown scars, coral bleaching and coral disease (black band disease, white syndrome, brown band disease, porites pinking, skeletal eroding band disease, hyperplasia and neoplasia).

Introduction to Reef Impacts and Coral Mortality

SCUBA searches have been used by the LTMP to provide information on sources of coral mortality, which assist in examining the reef in greater detail and interpreting trends in benthic cover at permanent sites. SCUBA searches enable:

- I. The detection of low-level populations of COTS. At low densities they are cryptic and more difficult to detect by methodologies such as the manta tow.
- II. SCUBA searches provide a method for the detection of juvenile COTS, which because of their small size and cryptic behaviour, are not easily seen in benthic or 3-Dimensional modelling assays.
- III. SCUBA searches enable the diver to detect other factors that may be causing coral mortality such as *Drupella* spp., bleaching or disease (e.g. white syndromes and black band disease).

4. RESULTS


4.1 Benthic Cover

Results of overall benthic cover at Poblacion reef show abiotic categories (rock, rubble and sand) to be the dominant substrate type (42.99%), this category being formed primarily of rubble. This was followed by coral (32.17%) and algae (11.59%) (fig. 4.1). A total of 29 Scleractinian coral genera were recorded, with *Echinopora* (12.50%), *Porites* (6.38%), and *Acropora* (2.85%), proving to be the most abundant coral genera (fig 1.2). The site recorded a Simpson’s Index 1-D of 0.70.

4.2 Reef Impacts & Coral Mortality


Coral bleaching was the most prevalent impact recorded within Poblacion reef, with 22 instances recorded, affecting on average 40.9% of the colony tissue. A total of 10 coral genera were affected by bleaching, with *Fungia spp.* recording the highest bleaching prevalence. The average percentage of affected tissue is 36.05%.

Five (5) counts of coral disease were present in the form of neoplasia, white syndrome and porites pinkening. Affected colonies were from the genus *Porites spp.* and *Fungia spp.* Two (2) counts of trash were recorded within the surveyed space. No *Acanthaster plancii* (Crown of Thorns Starfish) or suspected feeding scars were recorded. Three (3) counts of *Drupella spp.* feeding activity was recorded affecting *Fungia spp.*, *Pocillopora spp.*, and *Galaxea spp.* colonies respectively. On average, *Drupella spp.* affected 33.12% of the recorded colonies.



Major Benthic Categories

Figure 4.1 Average percentage cover of all major benthic categories with standard error (\pm SE) recorded at Poblacion reef during the dry season of February to July 2019.



Predominant Coral Genera

Figure 4.2 Average percentage cover of the five major coral genera with standard error (\pm SE) recorded at Poblacion reef during the dry season of February to July 2019.

Table 1. Reef impacts recorded at Poblacion reef during the dry season of February to July 2019 with ranking and trends.

Measurement	Current Value	Ranking	Last Season Value	Trend
Coral Bleaching (count/100m ²)	5.5	1 st	n/a	n/a
Disease (incidences/100m ²)	1.25	1 st	n/a	n/a
<i>Acanthaster plancii</i> (count/100m ²)	0	5 th	n/a	n/a
<i>Drupella spp.</i> (count/100m ²)	0.75	6 th	n/a	n/a
Trash (count/100m ²)	0	9 th	n/a	n/a

4.3 Fish

Poblacion reef recorded an average total fish abundance of $n = 559$ across the two survey depths, a species richness (S) of $S = 62$, and an average total biomass of $9.13 \text{ kg}/250\text{m}^2$. *Pomacentridae* (damselfish) had both the highest recorded average abundance and species richness ($n = 434$, $S = 17$), followed by *Labridae* (wrasse) ($n = 415$, $S = 14$), *Acanthuridae* (surgeonfish) ($n = 28$, $S = 5$), and *Serranidae* (grouper) ($n = 12$, $S = 1$).

Scaridae had the highest total biomass at $0.54 \text{ kg}/250\text{m}^2$, followed by *Mullidae* and *Labridae* with biomass totals of $0.33 \text{ kg}/250\text{m}^2$ and $0.24 \text{ kg}/250\text{m}^2$ respectively (figure 4.3). Grouping fish into trophic groups showed that on average, across depths, most fish were planktivores ($n = 294$, $S = 8$) followed by omnivores ($n = 160$, $S = 2.5$). The trophic group with the greatest average total biomass were the herbivores ($0.42 \text{ kg}/250\text{m}^2$) followed by invertivores ($0.31 \text{ kg}/250\text{m}^2$) and planktivores ($0.21 \text{ kg}/250\text{m}^2$). Fish that fit within two trophic groups were counted separately in each group. Those within three or more groups were counted as omnivores (figure 4.4).

Commercially important families recorded at Sahara reef include *Scaridae* ($0.54 \text{ kg}/250\text{m}^2$), *Mullidae* ($0.33 \text{ kg}/250\text{m}^2$), *Labridae* ($0.24 \text{ kg}/250\text{m}^2$), *Acanthuridae* ($0.21 \text{ kg}/250\text{m}^2$), *Lutjanidae* ($0.14 \text{ kg}/250\text{m}^2$) and *Serranidae* ($0.007 \text{ kg}/250\text{m}^2$), which can be seen in figure 4.5. No *Haemulidae* (sweetlips) or *Lethrinidae* (emperor) were found within the transect.




Figure 4.3 Average total biomass ($\text{kg}/250\text{m}^2$) of fish families recorded at Poblacion reef during the dry season of February to July 2019.




Figure 4.4 Average total biomass ($\text{kg}/250\text{m}^2$) of fish trophic groups recorded at Poblacion reef during the dry season of February to July 2019.




Figure 4.5 Average biomass ($\text{kg}/250\text{m}^2$) commercially important fish families recorded at Poblacion reef during the dry season of February to July 2019.

4.3 Reef Complexity

Results from our 3-Dimensional reef reconstructions reveal an average rugosity index of 7.13 ± 2.07 , and a slope value of 0.13 ± 0.06 . Figures 4.6 and 4.7 show the rugosity across the surveyed depths of Poblacion reef.

Table 2. Summary of findings recorded at Poblacion reef during the dry season of February to July 2019 with ranking and trends.

Measurement	Current Value	Ranking	Last Season Value	Trend
Coral Cover (%)	32	3 rd	n/a	n/a
Algal Cover (%)	11.5	1 st	n/a	n/a
Coral 1-D	0.70	5 th	n/a	n/a
No. of Fish (count/250kg ²)	559	2 nd	n/a	n/a
Fish Biomass (kg/250m ²)	9.12	4 th	n/a	n/a
Fish 1-D	0.73	11 th	n/a	n/a
Rugosity (RQ)	7.13	2 nd	n/a	n/a





Figure 4.6 a) Average slope along the transect at 5m (replicate B). Scale is in mega pixels with 300MP being equal to 50 meters on the transect. b) Average rugosity along the transect at 5m (replicate B). Scale is in mega pixels with 300MP being equal to 50 meters on the transect. b) Digital Elevation Model (DEM) produced with SfM photogrammetry techniques at 5m (replicate B).




Figure 4.6 a) Average slope along the transect at 10m (replicate B). Scale is in mega pixels with 300MP being equal to 50 meters on the transect. b) Average rugosity along the transect at 10m (replicate B). Scale is in mega pixels with 300MP being equal to 50 meters on the transect. b) Digital Elevation Model (DEM) produced with SfM photogrammetry techniques at 5m (replicate B).

5. DISCUSSION

This report documents the current status of the reef assemblage and community make-up of Poblacion reef, with findings contributing to the baseline dataset of the Dauin LTRMP.

Poblacion reef currently supports a relatively high percentage of both coral cover and diversity, ranked as 3rd and 5th within Dauin respectively. Examining the health of the recorded coral cover within Poblacion reef does reveal that a variety of stressors are compromising the health of select colonies within this reef system. Coral bleaching was present, with 22 cases across 10 different coral genera. This bleaching was recorded to be patchy across the colony, with an average of 41% of the colony tissue to be affected by the bleaching. Whether mortality follows bleaching in these circumstances is thought to be largely dependent on the coral's ability to endure starvation, or to supplement its energy requirements from food particles captured from the water column (heterotrophy)^{61,62,63}. Coral health prior to exposure to heat stress may be the most important factor influencing colony survivorship during bleaching events⁶¹. Most corals rely heavily on the energy produced by their zooxanthellae, which are expelled during a bleaching event⁶¹. As a result, corals in the bleached state are beginning to starve, and their ability to endure this hardship is likely to be an important determination for their survival^{61,62,63}. Like many animals, corals store surplus energy as lipids (fats). Corals in good condition will have relatively high lipid levels, providing a buffer against low energy supply. For this reason, it is thought that the condition of a coral at the time it bleaches may play a key role in determining whether it will be able to survive the period of starvation that follows^{61,62,63}. Whether mortality follows these bleaching cases at Poblacion reef

will continue to be monitored. Additional threats to coral within Poblacion reef include disease and low counts of predation by the corallivorous invertebrate *Drupella* spp. White syndrome disease is of the most concern, as this disease can cause rapid and irreversible damage. It is well established that infectious diseases are the outcome of interactions between susceptible hosts, disease agents, and the environment that live within⁶⁴. Semi-Permeable Membrane Devices (SPMDs) have been placed within Poblacion reef as a result of these findings. SPMDs are an effective way of monitoring the concentrations of chemicals from anthropogenic runoff and pollution in the marine environment due to their ability to detect miniscule levels of chemicals. Ongoing monitoring of this site is required to document the long-term ecological impacts of coral bleaching, disease and corallivorous activity. This will allow the assessment of the both the rate of recovery from these disturbances, as well as the cause of change to reef condition.

The species richness and abundance of reef fish is often associated with topographic or structural complexity^{35,65}. High topographic complexity may promote a high fish abundance as it provides more refuges, decreases the encounter rates between competitors, consequently reducing the effects of predation and competition^{35,65}. With Poblacion reef ranking 2nd for both rugosity and fish abundance, it is possible that this theory stands true for this reef system. However, more data will be required to test this correlation at Poblacion reef. Percent hard coral cover has also been identified as an important predictor of fish community structure, with fish species richness and abundance increasing with hard coral cover^{58,60}. With the current findings at Poblacion reef, it is expected that coral cover has had a greater influence on fish abundance than fish species richness. This may be

expected, due to a higher coral cover increasing the habitat area for coral-associated fish, while not necessarily increasing the range of available habitat types. It has been identified by Wilson *et al.* (2006) in a meta-analysis that fish declines becomes prominent in a coral cover decline of approximately 10%⁶⁰. Therefore it should be the change in coral cover at Poblacion reef, which will be outlined with future data collected as part of the Dauin LTRMP, that will be an important predictor of fish thresholds.

Additional studies support the factors of coral cover and coral diversity to be indicative of supporting a high fish diversity^{57,58,59}, however Poblacion reef has the lowest diversity of fish of all surveyed sites within the Dauin inshore reef. There are a number of reasons to expect a reduction of fish species richness due to coral diversity. Most reef fish have specific microhabitat requirements^{53,54}. Many of these requirements are highly dependent on a narrow suite of coral species or coral morphology for shelter or reproduction sites^{56,57}. However, because coral species differ greatly in their growth form and branching structure, it makes it difficult to assess the relative importance of different habitat variables in structuring fish communities. Messmer *et al.* (2011) recently demonstrated, by constructing small patch reefs with different coral assemblages, that reefs of higher diversity generally supported more species⁵⁷. However, these findings also revealed that the effects of coral diversity was explained by the inclusion of certain, highly preferred coral species⁵⁷. It is the loss of coral species preferred by the greatest range of fish species that will have the greatest impact on the fish assemblage⁵⁷. Further studies at Poblacion reef will be required to determine what coral species are favoured by a broad range of reef fishes, and if habitat specialisation would explain a coral species richness – fish species richness relationship.

The abundance and biomass of herbivorous fish outweighs that of coral-associated fishes at Poblacion reef. This presence is typically linked to habitat characteristics; for example algal presence. When delving into these findings, they do support the general expectation that a higher biomass of herbivorous fishes will be associated with lower levels of turf and fleshy macroalgal cover, and higher cover of reef calcifiers. Whilst *Poblacion* reef has the highest percentage of algal cover relative to the surrounding survey sites, a closer look reveals 4.3% of algal cover to be comprised of coralline algae, with the remaining 7.2% to comprise of macro and turf algae. Further analysis of herbivorous functional groups will be important to gain a broader understanding of the role herbivorous fish at Poblacion reef are playing in reef resilience.

These results hint towards current threats to Poblacion reef being as a result of climate change, land run-off, and perhaps synergistically coastal development. With Poblacion maintaining a MPA with daily supervision from Bantay Dagats, fishing within this reef should be absent. The scale at which these notified threats are prominent within Poblacion reef, the threshold at which the coral reef assemblage will respond, and the effectiveness of this MPA protection will be reported with ongoing assessment.

6. REFERENCES

1. Bell, S., McCoy, E., Mushinsky, H (1991). Habitat structure; the physical arrangement of objects in space, Bury St Edmunds, Suffolk, UK: St Edmundsbury Press Ltd.
2. Ferrigno, F., Bianchi, C., Lasagna, R., Morri, C., Russo, G., Sandulli, R. (2016). Corals in high diversity reefs resist human impact, *Ecological Indicators* 70, pp. 106-113
4. Komyakova, V., Munday, P.L., Jones, G.P. (2013). Relative Importance of Coral Cover, Habitat Complexity and Diversity in Determining the Structure of Reef Fish Communities. *PLOS ONE* 8(12): e83178.
5. Lirman, D., Fong, P. (1997). Patterns of Damage to the Branching Coral *Acropora palmata* Following Hurricane Andrew: Damage and Survivorship of Hurricane-Generated Asexual Recruits, *Journal of Coastal Research* 13(1) pp. 67-72
6. Quimpo, T.J.R., Cabaitan, P.C., Go, K.T.B., Dumalagan, E.E. Jr, Villanoy, C.L., Siringan, F.P. (2019). Similarity in benthic habitat and fish assemblages in the upper mesophotic and shallow water reefs in the West Philippine Sea. *Journal of the Marine Biological Association of the United Kingdom* 1(11)
7. Roff, G., Mumby, P. (2012). Global disparity in the resilience of coral reefs, *Trends in ecology and evolution* 27(7), pp 404-417,
8. Bohnsack, J.A., Harper, D.E., McClellan, D.B., Hulsbeck, M. (1994). Effects of reef size on colonization and assemblage structure of fishes at artificial reefs off southeastern Florida, USA. *Bull Mar Sci.* 55(2-3):796-823.
9. Burt, J., Bartholomew, A., Bauman, A., Saif, A., Sale, P.F. (2009). Coral recruitment and early benthic community development on several materials used in the construction of artificial reefs and breakwaters. *J Exp Mar Bio Ecol.* 373(1):72-8.
10. Baynes, T.W., Szmant, A.M. (1989). Effect of current on the sessile benthic community structure of an artificial reef. *Bull Mar Sci.* 144(2):545-66.
11. Brown, C.J. (2005). Epifaunal colonization of the Loch Linnhe artificial reef: influence of substratum on epifaunal assemblage structure. *Biofouling.* 21(2):73-85. pmid:16167388
12. Andradi-Brown, D.A., Gress, E., Wright, G., Exton, D.A., Rogers, A.D. (2016). Reef fish community biomass and trophic structure changes across shallow to upper-mesophotic reefs in the Mesoamerican Barrier Reef, Caribbean. *PLoS ONE* 11(6): e0156641.
13. Bianchessi, A (2012). A Case Study of the Municipality of Dauin, Negros Oriental, Philippines: A Global Bright Spot for Marine Conservation. University of Texas, El Paso.
14. Burns, J., Delparte, D., Gates, R., Takabayashi, M. (2015). Integrating Structure-from-Motion Photogrammetry with Geospatial Software as a Novel Technique for Quantifying 3D Ecological Characteristics of Coral Reefs. *PeerJ.*
15. Burrough, P.A., McDonell, R.A. (1998). Principles of geographical information systems. New York: Oxford University Press. 190 pp.

16. Chun, J.B., Jung, H., Kyung, C.M. (2008). Suppressing rolling-shutter distortion of CMOS image sensors by motion vector detection. *IEEE Transactions on Consumer Electronics* 54(4):1479–1487.
17. De’ath, G., Fabricius, K.E. (2000). Classification and regression trees: A powerful yet simple technique for ecological data analysis. *Ecology*, 81(11), pp. 3178-3192.
18. De’ath, G., Fabricius, K.E., Sweatman, H., Poutinen, M. (2012). The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Australian Institute of Marine Science*. 109(44), pp. 17995-17999.
19. De Smith, M.J., Goodchild, M.F., Longley, P.A. (2015). *Geospatial analysis: a comprehensive guide to principles, techniques, and software tools*. Fifth edition, Issue version: 1. Alberta: Winchelsea Press.
20. Figueira, W., Ferrari, R., Weatherby, E., Porter, A., Hawes, S., Byrne, M (2015). Accuracy and Precision of Habitat Structural Complexity Metrics Derived from Underwater Photogrammetry. *Remote Sens.* 7, 16883-16900.
21. Friedlander, A.M., Parrish, J.D. (1998). Habitat characteristics affecting fish assemblages on a Hawaiian coral reef. *Journal of Experimental Marine Biology and Ecology* 224:1–30
22. Great Barrier Reef Marine Park Authority 2014, Great Barrier Reef Outlook Report 2014, GBRMPA, Townsville.
23. Hoegh-Guldberg, O., Hoegh-Guldberg, H., Veron, J.E.N., Green, A., Gomez, E. D., Lough, J., King, M., Ambaryanto, Hansen, L., Cinner, J., Dews, G., Russ, G., Schuttenberg, H. Z., Peñaflores, E.L., Eakin, C. M., Christensen, T. R. L., Abbey, M., Areki, F., Kosaka, R. A., Tewfik, A., Oliver, J. (2009). *The Coral Triangle and Climate Change: Ecosystems, People and Societies at Risk*. WWF Australia, Brisbane, 276 pp.
24. Jonker, M. (Michelle) II. Australian Institute of Marine Science III. Title (Series: Long- term monitoring of the Great Barrier Reef standard operational procedure; 10).
25. Komyakova, V., Munday, P.L., Jones, G.P. (2013). Relative importance of coral cover, habitat complexity and diversity in determining the structure of reef fish communities. *PLoS ONE*, 8(12), e83178.
26. Kostylev, V.E., Erlandsson, J., Ming, M.Y., Williams, G.A. (2005). The relative importance of habitat complexity and surface area in assessing biodiversity: fractal application on rocky shores. *Ecological Complexity* 2:272–286
27. Luckhurst, B.E., Luckhurst, K. (1978). Analysis of the influence of substrate variables on coral reef fish communities. *Marine Biology* 49:317–323
28. McCormick, M. (1994). Comparison of field methods for measuring surface topography and their associations with a tropical reef fish assemblage. *Marine Ecology Progress Series* 112:87–96
29. Miller, I. R. (2003) Crown-of-thorns starfish and coral surveys using the manta tow and scuba search techniques. Long-term Monitoring of the Great Barrier Reef. Standard Operational Procedure, No. 8, Australian Institute of Marine Science, Townsville, 43pp.

30. Pittman, S.J., Brown, K.A. (2011). Multi-scale approach for predicting fish species distributions across coral reef seascapes. *PLoS ONE* 6(5):e20583
31. Pittman, S.J., Costa, B.M., Battista, T.A. (2009). Using lidar bathymetry and boosted regression trees to predict the diversity and abundance of fish and corals. *Journal of Coastal Research* 25(6):27–38
32. Raoult et al. (2016). GoPros™ as an underwater photogrammetry tool for citizen science. *PeerJ* 4:e1960
33. Sweatman, H., Burgess, S., Cheal, A., Coleman, G., Delean, S., Emslie, M., McDonald, A., Miller, I., Osborne, K., and Thompson, A. (2005). Long-term Monitoring of the Great Barrier Reef Status Report, No. 7, Australian Institute of Marine Science, Townsville, 266 pp.
34. Verhoeven, G. (2012). Getting computer vision airborne: using structure from motion for accurate orthophoto production. In: RSPSoc archaeology special interest group meeting spring 2012. 4–6
35. Young, G.C., Dey, S., Rogers, A.D., Exton, D. (2018). Correction: Cost and time-effective method for multi-scale measures of rugosity, fractal dimension, and vector dispersion from coral reef 3D models. *PLOS ONE* 13(7): e0201847.
36. Connell, J.H., Slatyer, R.O. (1977). Mechanisms of succession in natural communities and their role in community stability and organization. *Am Nat.* 111(982):1119–44.
37. Schuhmacher, H. (1988). Development of coral communities on artificial reef types over 20 years (Eilat, Red Sea). In *Proc 6th Coral Reef Symp.* (3):379–384.
38. Bailey-Brock, J.H. (1989). Fouling community development on an artificial reef in Hawaiian waters. *Bull Mar Sci.* 44(2):580–91.
39. Cummings, S.L. (1994). Colonization of a nearshore artificial reef at Boca Raton (Palm Beach County), Florida. *Bull Mar Sci* 55:1193-1215
40. Palmer-Zwahlen, M.L., Aseltine, D.A. (1994). Successional development of the turf community on a quarry rock artificial reef. *Bull Mar Sci* 55:920-923
41. Burke, Laretta & Selig, Elizabeth & Spalding, Mark & authors, other & McManus, John. (2002). *Reefs at Risk in South East Asia.*
42. Allen, G. R. (2007). Conservation hotspots of biodiversity and endemism for Indo-Pacific coral reef fishes. *Aquatic Conserv: Mar. Freshw. Ecosyst.* DOI: 10.1002/aqc.880.
43. Snavely, N., Seitz, S.N., Szeliski, R. (2008). Modelling the world from internet photo collections. *International Journal of Computer Vision.* 80:189–210. doi: 10.1007/s11263-007-0107-3.
44. Westoby, M.J., Brasington, J., Glasser, N.F., Hambrey, M.J., Reynolds, J.M. (2012). ‘Structure-from-Motion’ photogrammetry: a low-cost, effective tool for geoscience applications. *Geomorphology.* 2012;179:300–314.
45. Fisher, W.S., Davis, W.P., Quarles, R.L., Patrick, J., Campbell, J.G., Harris, P.S., Hemmer, B.L., Parsons, M. (2007). Characterizing coral condition using estimates of three-dimensional colony surface area. *Environmental*

- Monitoring and Assessment. 125:347–360.
46. Done TJ. (1997). Decadal changes in reef-building communities: implications for reef growth and monitoring programs. Proceedings of the 8th International Coral Reef Symposium. 1:411–416.
 47. Figueira W, Ferrari R, Weatherby E, Porter A, Hawes S, Byrne M (2015). Accuracy and Precision of Habitat Structural Complexity Metrics Derived from Underwater Photogrammetry. Remote Sensing. 7(12):16883–16900.
 48. Holmes, T.H., Wilson, S.K., Travers, M.J., Langlois, T.J., Evans, R.D., Moore, G.I., Douglas, R.A., Shedrawi, G., Harvey, E.S., and Hickey, K. (2013). A comparison of visual-and stereo-video based fish community assessment methods in tropical and temperate marine waters of Western Australia. Limnology and oceanography, methods. 11(7), pp.337-350.
 49. Stopnitzky S. (2014). Studying Reefs, Staying Dry. Khaleb bin Sultan Living Oceans Foundation. <https://www.livingoceansfoundation.org/studying-reefs-staying-dry/>
 50. Kohler, K.E., Gill, S.M. (2006). Coral Point Count with Excel extensions (CPCe): A Visual Basic program for the determination of coral and substrate coverage using random point count methodology. Computers & Geosciences, 32: 1259-1269
 51. Rogers, C.S., Garrison, G., Grober, R., Hillis, Z.M., Franke, M.A. (1994). Coral reef monitoring manual for the Caribbean and Western Atlantic. Virgin Islands National Park, St. John, US Virgin Islands, 107, USA
 52. Rogers, C.S. (1988). Recommendations for long-term assessment of coral reefs: US National Park initiates regional program. Proc 6th Int. Coral Reef Symposium, Australia, 2: 339-403.
 53. Depczynski, M., Bellwood, D.R. (2004) Microhabitat utilisation patterns in cryptobenthic coral reef fish communities. Marine Biology 145: 455-463.
 54. Munday PL, Jones GP (1998) The ecological implications of small body size among coral-reef fishes. Oceanogr Mar Biol Annu Rev 36: 373-411.
 55. Habitat preferences of the fishes that dwell in shrub corals on the Great Barrier Reef. Proc Acad Nat Sci Phila 123: 1-26.
 56. Sale, P.F. (1977) Maintenance of high diversity in coral reef fish communities. Am Nat 111: 337-359.
 57. Messmer, V., Jones, G.P., Munday, P.L., Holbrook, S.J., Schmitt, R.J. (2011) Habitat biodiversity as a determinant of fish community structure on coral reefs. Ecology 92(12): 2285-2298
 58. Chabanet, P., Ralambondrainy, H., Amanieu, M., Faure, G., Galzine, R. (1997) Relationships between coral reef substrata and fish. Coral Reefs 16: 93-102.
 59. Jones, G.P., McCormick, M.I., Srinivasan, M., Eagle, J.V. (2004) Coral decline threatens fish biodiversity. Proc Natl Acad Sci U S A 101: 8251-8253.
 60. Wilson, S.K., Dolman, A.M., Cheal, A.J., Emslie, M.J., Pratchett, M.S. (2009) Maintenance of fish diversity on disturbed coral reefs. Coral Reefs 28: 3-14.

61. Hoegh-Guldberg O (1999). Climate change, coral bleaching and the future of the world's coral reefs. *Marine and Freshwater Research* 50:839–866.
62. Muscatine, L. (1990). The role of symbiotic algae in carbon and energy flux in coral reefs. *Coral Reefs* 25:75–87.
63. Brown, B.E. (1997). Coral bleaching: causes and consequences. *Coral Reefs* 16:129–S138.
64. Bruckner, A.W. (2002). Priorities for effective management of coral diseases. NMFS-OPR-22. National Oceanic and Atmospheric Administration, Silver Spring, MD
65. Wilson, S.K., Graham, N.A.J., Polunin, N.V.C. (2007) Appraisal of visual assessments of habitat complexity and benthic composition on coral reefs. *Mar Biol* 151: 1069-1076.